



THE UNIVERSITY of EDINBURGH

Edinburgh Research Explorer

Life-Course Genome-wide Association Study Meta-analysis of Total Body BMD and Assessment of Age-Specific Effects

Citation for published version:

Medina-Gomez, C, Kemp, JP, Trajanoska, K, Luan, J, Chesi, A, Ahluwalia, TS, Mook-Kanamori, DO, Ham, A, Hartwig, FP, Evans, DS, Joro, R, Nedeljkovic, I, Zheng, H-F, Zhu, K, Atalay, M, Liu, C-T, Nethander, M, Broer, L, Porleifsson, G, Mullin, BH, Handelman, SK, Nalls, MA, Jessen, LE, Heppe, DHM, Richards, JB, Wang, C, Chawes, B, Schraut, KE, Amin, N, Wareham, N, Karasik, D, Van der Velde, N, Ikram, MA, Zemel, BS, Zhou, Y, Carlsson, CJ, Liu, Y, McGuigan, FE, Boer, CG, Bønnelykke, K, Ralston, SH, Robbins, JA, Walsh, JP, Zillikens, MC, Langenberg, C, Li-Gao, R, Williams, FMK, Harris, TB, Akesson, K, Jackson, RD, Sigurdsson, G, den Heijer, M, van der Eerden, BCJ, van de Peppel, J, Spector, TD, Pennell, C, Horta, BL, Felix, JF, Zhao, JH, Wilson, SG, de Mutsert, R, Bisgaard, H, Styrkarsdóttir, U, Jaddoe, VW, Orwoll, E, Lakka, TA, Scott, R, Grant, SFA, Lorentzon, M, van Duijn, CM, Wilson, JF, Stefansson, K, Psaty, BM, Kiel, DP, Ohlsson, C, Ntzani, E, van Wijnen, AJ, Forgetta, V, Ghanbari, M, Logan, JG, Williams, GR, Bassett, JHD, Croucher, PI, Evangelou, E, Uitterlinden, AG, Ackert-Bicknell, CL, Tobias, JH, Evans, DM & Rivadeneira, F 2018, 'Life-Course Genome-wide Association Study Meta-analysis of Total Body BMD and Assessment of Age-Specific Effects', *American Journal of Human Genetics*, vol. 102, no. 1, pp. 88-102. <https://doi.org/10.1016/j.ajhg.2017.12.005>

Digital Object Identifier (DOI):

[10.1016/j.ajhg.2017.12.005](https://doi.org/10.1016/j.ajhg.2017.12.005)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

American Journal of Human Genetics

Publisher Rights Statement:

This is the author's peer-reviewed manuscript as accepted for publication.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 08. May. 2023



Life-course Genome-Wide Association Study Meta-analysis of Total Body BMD and

Assessment of Age-specific Effects

Carolina Medina-Gomez^{1,2,3,63}, John P. Kemp^{4,5,63}, Katerina Trajanoska^{1,3}, Jian'an Luan⁶,
Alessandra Chesi⁷, Tarunveer S. Ahluwalia^{8,9}, Dennis O. Mook-Kanamori^{10,11}, Annelies Ham¹,
Fernando P. Hartwig¹², Daniel S. Evans¹³, Raimo Joro¹⁴, Ivana Nedeljkovic³, Hou-Feng
Zheng^{15,16,17}, Kun Zhu^{18,19}, Mustafa Atalay¹⁴, Ching-Ti Liu²⁰, Maria Nethander²¹, Linda Broer¹,
Gudmar Porleifsson²², Benjamin H. Mullin^{18,19}, Samuel K. Handelman²³, Mike A. Nalls^{24,25}, Leon
E. Jessen⁸, Denise H.M. Heppel^{2,3}, J. Brent Richards^{15,16}, Carol Wang²⁶, Bo Chawes⁸, Katharina E
Schraut^{27,28}, Najaf Amin², Nick Wareham⁶, David Karasik^{29,30}, Nathalie Van der Velde^{1,31}, M.
Arfan Ikram³, Babette S. Zemel³², Yanhua Zhou²⁰, Christian J. Carlsson⁸, Yongmei Liu³³, Fiona E.
McGuigan³⁴, Cindy G. Boer¹, Klaus Bønnelykke⁸, Stuart H Ralston³⁵, John A. Robbins³⁶, John P.
Walsh^{18,19}, M. Carola Zillikens¹, Claudia Langenberg⁶, Ruifang Li-Gao¹⁰, Frances M.K. Williams³⁷,
Tamara B. Harris³⁸, Kristina Akesson^{34,39}, Rebecca D. Jackson⁴⁰, Gunnar Sigurdsson^{41,42}, Martin
den Heijer^{10,43}, Bram C.J. van der Eerden¹, Jeroen van de Peppel¹, Timothy D. Spector³⁷, Craig
Pennell²⁶, Bernardo L Horta¹², Janine F. Felix^{2,3,44}, Jing Hua Zhao⁶, Scott G. Wilson^{18,19,37}, Renée
de Mutsert¹⁰, Hans Bisgaard⁸, Unnur Styrkarsdóttir²², Vincent W. Jaddoe^{2,3,44}, Eric Orwoll⁴⁵,
Timo A. Lakka^{14,46,47}, Robert Scott⁶, Struan F.A. Grant^{7,48}, Mattias Lorentzon⁴⁹, Cornelia M van
Duijn³, James F. Wilson^{27,50}, Kari Stefansson^{22,41}, Bruce M Psaty^{51,52}, Douglas P Kiel^{28,53,54}, Claes
Ohlsson⁵⁵, Andre J. van Wijnen⁵⁷, Vincenzo Forgetta^{15,16}, Mohsen Ghanbari^{3,58}, Graham R.
Williams⁵⁹, J.H. Duncan Bassett⁵⁹, Evangelos Evangelou^{56,60}, Andre G. Uitterlinden^{1,2,3}, Cheryl L
Ackert-Bicknell⁶¹, Jonathan H. Tobias⁶², David M. Evans^{4,5,64}, Fernando Rivadeneira^{1,2,3,64}

1) Department of Internal Medicine, Erasmus MC, 3000 CA Rotterdam, The Netherlands; 2) The Generation R Study Group, Erasmus MC, 3000 CA Rotterdam, The Netherlands; 3) Department of Epidemiology, Erasmus MC, 3000 CA Rotterdam, The Netherlands; 4) University of Queensland Diamantina Institute, Translational Research Institute, Brisbane, Queensland 4102, Australia; 5) MRC Integrative Epidemiology Unit, University of Bristol, Bristol BS8 2BN, UK; 6) MRC Epidemiology Unit, University of Cambridge, Cambridge CB2 0QQ, UK; 7) Division of Human Genetics, Children's Hospital of Philadelphia, Philadelphia, PA 19104, USA; 8) COPSAC, Copenhagen Prospective Studies on Asthma in Childhood, Herlev and Gentofte Hospital, University of Copenhagen, Copenhagen 2820, Denmark; 9) Steno Diabetes Center Copenhagen 2820, Denmark; 10) Department of Clinical Epidemiology, Leiden University Medical Centre, 2333 ZA Leiden, The Netherlands; 11) Department of Public Health and Primary Care, Leiden University Medical Centre, 2333 ZA Leiden, The Netherlands; 12) Postgraduate Program in Epidemiology, Federal University of Pelotas, 96020220 Pelotas, Brazil; 13) California Pacific Medical Center Research Institute, San Francisco, CA 94107, USA; 14) Institute of Biomedicine, Physiology, University of Eastern Finland, Kuopio 70211, Finland; 15) Departments of Medicine, Human Genetics, Epidemiology and Biostatistics, McGill University, Montreal H3A 1A2, Canada; 16) Department of Medicine, Lady Davis Institute for Medical Research, Jewish General Hospital, McGill University, Montreal H3T 1E2, Canada; 17) Hangzhou Normal University, 311121 Zhejiang, China; 18) Department of Endocrinology and Diabetes, Sir Charles Gairdner Hospital, Nedlands 6009, Australia; 19) School of Medicine and Pharmacology, University of Western Australia, Crawley 6009, Australia; 20) Department of Biostatistics, Boston University School of Public Health, Boston, MA 02131, USA; 21) Bioinformatics Core Facility, Sahlgrenska Academy, University of Gothenburg, 41390 Gothenburg, Sweden; 22) deCODE Genetics/Amgen, Reykjavik, IC-101, Iceland; 23) Department of Obstetrics and Gynecology, Wayne State University, Detroit, MI 48201, USA; 24) Laboratory of Neurogenetics, National Institute on Aging, Bethesda, MD 20892, USA; 25) Data Tecnica International, Glen Echo, MD 20812, USA; 26) School of Women's and Infants' Health, University of Western Australia, Crawley 6009, Australia; 27) Centre for Global Health Research, Usher Institute for Population Health Sciences and Informatics, University of Edinburgh, Edinburgh EH16 4UX, Scotland; 28) Centre for Cardiovascular Sciences, Queen's Medical Research

Institute, University of Edinburgh, Edinburgh EH16 4TJ, Scotland; 29) Hebrew SeniorLife, Institute for Aging Research, Roslindale, MA 02131, USA; 30) Faculty of Medicine in the Galilee, Bar-Ilan University, Safed 1311502, Israel; 31) Department of Internal Medicine, Section of Geriatrics, Academic Medical Center, 1105 AZ Amsterdam, The Netherlands; 32) Division of GI, Hepatology, and Nutrition, Children's Hospital of Philadelphia, Philadelphia, PA 19104, USA; 33) Wake Forest School of Medicine Winston-Salem, NC 27101, USA; 34) Lund University, Department of Clinical Sciences Malmö, Clinical and Molecular Osteoporosis Research Unit, 20502 Malmö, Sweden; 35) Centre for Genomic and Experimental Medicine, Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh EH4 2XU, Scotland; 36) Department of Medicine, University of California, Davis; Sacramento, CA 95817, USA; 37) Department of Twin Research & Genetic Epidemiology, King's College London, London SE1 7EH, UK; 38) Laboratory of Epidemiology and Population Sciences, National Institute on Aging, Bethesda, MD 20892, USA; 39) Skåne University Hospital, Department of Orthopedics, 21428 Malmö, Sweden; 40) Division of Endocrinology, Diabetes, and Metabolism, Department of Internal Medicine, The Ohio State University, Columbus, OH 43210, USA; 41) Department of Endocrinology and Metabolism, Landspítali, The National University Hospital of Iceland, Reykjavik, IC-101, Iceland; 42) Faculty of Medicine, University of Iceland, Reykjavik, IC-101, Iceland; 43) Department of Clinical Epidemiology and Department of Internal Medicine, VU Medical Center, 1081 HV Amsterdam, The Netherlands; 44) Department of Pediatrics, Erasmus MC, 3000 CA Rotterdam, The Netherlands; 45) Department of Public Health & Preventive Medicine, Oregon Health & Science University, Oregon, Portland, OR 97239, USA; 46) Kuopio Research Institute of Exercise Medicine, Kuopio, 70100, Finland; 47) Department of Clinical Physiology and Nuclear Medicine, Kuopio University Hospital, Kuopio, 70210, Finland; 48) Department of Pediatrics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA; 49) Geriatric Medicine, Department of Internal Medicine and Clinical Nutrition, Institute of Medicine, University of Gothenburg, 40530 Gothenburg, Sweden; 50) MRC Human Genetics Unit, Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh EH4 2XU, Scotland; 51) Cardiovascular Health Research Unit, Departments of Medicine, Epidemiology, and Health Services, University of Washington, Seattle, WA 98101, USA; 52) Kaiser Permanente Washington Health Research Institute, Seattle, WA 98101, USA; 53) Department of Medicine Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, MA 02215, USA; 54) Broad Institute of MIT and Harvard, Boston, MA 02115, USA; 55) Center for Bone and Arthritis Research, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, 41345 Gothenburg, Sweden; 56) Department of Hygiene and Epidemiology, University of Ioannina Medical School, Ioannina, 45110, Greece; 57) Department of Orthopedic Surgery, Mayo Clinic, Rochester, MN 55905, USA; 58) Department of Genetics, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran; 59) Molecular Endocrinology Laboratory, Department of Medicine, Imperial College London, London W12 0NN, UK; 60) Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London W2 1PG, UK. 61) Center for Musculoskeletal Research, University of Rochester, Rochester, NY 14642, USA; 62) School of Clinical Sciences, University of Bristol, Bristol BS10 5NB, UK.

63)These authors contributed equally

64)These authors contributed equally

Corresponding author:

Dr. Fernando Rivadeneira MD PhD
Associate Professor Musculoskeletal Genomics
Genetic Laboratory - Department of Internal Medicine
Erasmus University Medical Center Rotterdam
The Netherlands
Email: f.rivadeneira@erasmusmc.nl

Abstract

Bone mineral density (BMD) assessed by DXA is used to evaluate bone health. In children, total body (TB) measurements are commonly used; in older individuals, BMD at the lumbar spine (LS) and femoral neck (FN) is used to diagnose osteoporosis. To date, genetic variants in more than 60 loci have been identified as associated with BMD. To investigate the genetic determinants of TB-BMD variation along the life course and test for age-specific effects, we performed a meta-analysis of 30 genome-wide association studies (GWAS) of TB-BMD including 66,628 individuals overall and divided across five age-strata each spanning 15 years. We identified variants associated with TB-BMD at 80 loci, of which 36 have not been previously identified; overall they explain approximately 10% of the TB-BMD variance when combining all age groups and influence the risk of fracture. Pathway and enrichment analysis of the association signals showed clustering within gene-sets implicated in the regulation of cell growth and SMAD proteins; overexpressed in the musculoskeletal system; and enrichment in enhancer and promoter regions. These findings reveal TB-BMD as a relevant trait for genetic studies of osteoporosis, enabling the identification of variants and pathways influencing different bone compartments. Only variants in *ESR1* and close proximity to *RANKL* showed a clear effect dependency on age. This most likely indicate that the majority of genetic variants identified influence BMD early in life and their effect can be captured throughout the life course.

Introduction

Osteoporosis is a disease characterized by low bone mass and microarchitectural deterioration of bone tissue leading to increased risk of fracture¹. It is diagnosed through the measurement

of bone mineral density (BMD) utilizing dual-energy X-ray absorptiometry (DXA), which is the single best predictor of fracture¹.

Bone is a dynamic tissue constantly undergoing resorption and formation. Bone mass increases steadily during childhood and markedly during adolescent growth². Peak bone mass is attained at approximately the third decade of life. Thereafter, until about 50 years of age, BMD remains fairly stable, by virtue of the coupling between bone formation and resorption (e.g., bone remodeling). Subsequently, bone resorption exceeds the rate of bone formation, resulting in a decrease in BMD, particularly in women after the onset of menopause³.

The International Society for Clinical Densitometry recommends performing DXA measurements at the lumbar spine, femoral neck and total hip to diagnose osteoporosis in postmenopausal women and men who are 50 years or older⁴. Consequently, studies of BMD determinants are frequently based on measurements at these skeletal sites. By contrast, for the assessment of bone health in children and adolescents, total body (excluding head) and lumbar spine are the preferred sites to minimize measurement artifacts resulting from changing areas in growing bones⁴. Nevertheless, in elderly individuals degenerative changes in the spine can give elevated BMD readings⁵. Moreover, total body DXA scans have been obtained in a number of adult research cohorts, primarily to assess body composition. Therefore, the total body BMD (TB-BMD) measurement is the most appropriate method for an unbiased assessment of BMD variation in the same skeletal site from childhood to old age.

To date, nearly 80 independent genetic variants have been shown to be robustly associated with variability in bone parameters⁶⁻¹⁸. Most of these markers have been identified in studies

comprising tens of thousands of adult and elderly individuals with DXA-derived BMD measurements, although a few of them have been associated with BMD specifically in studies of pediatric cohorts⁸. Furthermore, several of the associated variants display significant site-specific effects, possibly reflecting differences in bone composition across skeletal sites (e.g., cortical bone vs. trabecular bone) or differential response to mechanical loading⁸. Moreover, genetic studies on measures from peripheral quantitative computed tomography (pQCT) and bone quantitative ultrasound, which provide additional information regarding bone size, geometry and (micro) architecture identified genetic variants that may have specific effects on bone properties that are poorly captured by conventional DXA measurements⁹⁻¹⁰.

Given the complex physiological processes underlying age-related changes in BMD across the life course, it is possible that genetic studies in more refined age groups will reveal variants in unreported loci as well as age-specific genetic effects. Thus, the purpose of this study was to identify gene variants associated with TB-BMD across the life span and investigate possible differences of genetic effects across age periods.

Methods

TB-BMD GWAS meta-analyses

Study Populations

Subjects

This study comprised 30 epidemiological studies comprising ~66,628 individuals from populations across America, Europe, and Australia, with a variety of designs (**Supplemental Data; Table S1**) and participant characteristics (**Table S2**). In summary, most participants came from population-based cohorts of European ancestry (86%), two cohorts comprising African-American individuals (2%) and other four studies holding a fraction of individuals from admixed background (14%). All research aims and the specific measurements have been approved by the correspondent Medical Ethical Committee of each participating study. Written informed consent was provided by all subjects or their parents in the case of children.

BMD measurement

Total body BMD (g/cm^2) was measured by DXA following standard manufacturer protocols. As recommended by the International Society for Clinical Densitometry total body less head (TBLH) was the measurement used in pediatric cohorts⁴ (e.g., 0-15 years). Detailed information on the assessments performed by each study can be found in **Table S1**.

GWAS data and imputation

All individuals included in this study had genome-wide array data. Quality control of genotypes is summarized in **Table S1**. To enable meta-analysis, each study performed genotype

imputation using the cosmopolitan (all ethnicities combined) 1000 genomes phase 1 version 3 (March 2012) reference panel, yielding ~ 30,000,000 SNPs for analysis. Three studies used the combined 1000 genomes and the UK10K reference panels as presented in **Table S1**.

Association Analysis

TB(LH)-BMD was corrected for age, weight, height and genomic principal components (derived from GWAS data), as well as any additional study-specific covariates (e.g. recruiting center), in a linear regression model. For studies with non-related individuals, residuals were computed separately by sex, whereas for family-based studies sex was included as a covariate in the model. Finally, residuals were inverse normal transformed. The analyses were performed in each study for the overall population as well as in subgroups of individuals by age-strata, defined by bins of 15 years (i.e., 0-15 years, 15-30 years, 30-45 years, 45-60 years, and 60 or more years). SNP association was tested for autosomal variants, in which the additive effect of each SNP on the normalized BMD-residuals was estimated via linear regression.

Quality control of TB-BMD association summary statistics

A centralized quality-control procedure implemented in EasyQC¹⁹ was applied to all study-specific files of association results to identify cohort-specific issues. We excluded variants if they had missing information (e.g., missing association P-value, beta estimate, alleles, allele frequency), or nonsensical values (e.g., absolute beta estimates or standard errors >10, association P-values >1 or <0; or imputation quality < 0; infinite beta estimates or standard errors); minor allele frequency (MAF) less than 0.5%; imputation quality scores <0.4 (Impute2) or <0.3 (Minimac). Moreover, variants were flagged if they had large allele frequency

deviations from reference populations (>0.6 for admixed studies and >0.3 for ancestry-homogeneous studies).

GWAS meta-analyses

In the first instance, no exclusion criteria based on ancestry were applied for the meta-analysis ($N=66,628$). In addition, meta-analyses were carried out across age strata (minimum sample size per bin $N=200$ for each study) comprising: 1) 0-15 years ($N=11,807$), 15-30 years ($N=4,180$), 30-45 years ($N=10,062$), 45-60 years ($N=18,805$), and 60 or more years ($N=22,504$). Further, summary data from cohorts of European ancestry only were meta-analyzed and used in subsequent analyses. We discarded variants present in less than three studies. Approximately 23,700,000 markers (including SNPs and INDELS) were assessed for association. We applied the conventional genome-wide significance level (GWS, $P < 5 \times 10^{-8}$) for SNP discovery.

Assessment of Age-dependent effects

We selected SNPs which were suggestively (12,567 SNPs, $P < 5 \times 10^{-6}$) associated with BMD in the overall meta-analysis, present in at least 2 studies per age-bin and with MAF differences across these meta-analyses lower than 0.5. We clumped this dataset with an $r^2 \geq 0.8$, using as reference the most strongly associated SNPs with BMD and, pruning remaining SNPs within 0.7 Mb of each other. Age-dependent effects were assessed using a meta-regression approach for 1,464 SNPs obtained after this selection procedure. We ran a linear regression of the SNP effect estimates onto an intercept and the median age of each subgroup (e.g., each study stratified in age-bins). As proposed previously²⁰, standard errors of the effect estimates of each subgroup were multiplied by the square root of the genomic inflation factor when it was greater than 1.

We performed the meta-regression using the Metafor package²¹, and any statistical evidence of linear association was corrected for multiple testing (Bonferroni correction; $0.05/1,464 = 3.4 \times 10^{-5}$). The difference between beta-estimates in children vs. elderly meta-analyses (Pdiff) was tested using Easy-strata²².

Approximate conditional meta-analyses

Conditional analyses were undertaken based on the meta-analysis of the studies of European ancestry only (N=56,284). Only variants in the loci that reached GWS in this meta-analysis were assessed. The Rotterdam Study I (n=6,291) was used as reference for precise calculation of the linkage disequilibrium (LD) between the analyzed markers. We used an iterative strategy as implemented in GCTA²³ to determine: 1) independence of association signals within loci discovered in our study, by means of stepwise model selection procedure per chromosome (--massoc-slct routine); and 2) the novelty of the association signals discovered by our meta-analysis with regard to variants reported in previous well-powered GWAS of different bone traits (**Table S3**). To this end, we performed the association analysis conditional on 78 variants present in our data and associated with different bone-traits (--massoc-cond routine). These 78 SNPs were selected from different GWAS publications^{6-10;12-14}, assuring their independence to avoid collinearity issues.

Shared Genetic architecture of TB-BMD fracture and other traits

LD score regression analyses

We used the LD score regression package to estimate the heritability of TB-BMD and rule out that our results were a product of bias (e.g., residual population stratification or cryptic relatedness). LD score regression uses GWAS summary statistics and assesses the SNP-heritability based on the expected relationship between linkage disequilibrium (LD) of neighboring SNPs and strength of association under a polygenic model²⁴. As this methodology relies on the LD structure throughout the genome, we restricted this analysis to summary statistics from the meta-analysis of cohorts comprising only individuals from European ancestry. We used the publicly available, pre-computed LD structure data files specific to European populations of the HAPMAP 3 reference panel. An extension of this method allows estimating the genetic correlation between two traits²⁵. This can be performed in the LDhub pipeline, a web utility which gathers data from many different GWAS meta-analysis²⁶. From the 199 traits, currently available in the website, we have restricted our analysis to those traits whose heritability z-scores were larger than 4 and were analyzed only in European ancestry individuals (following the recommendations in the LD score software website (**Web Resources**)). Additionally, we incorporated data from a recent GWAS meta-analysis of any-type of fracture in individuals from European ancestry (N= 264,267; 37,778 cases) (K.T, unpublished data). In total, we assessed the genetic correlation between TB-BMD and 74 traits.

Mendelian randomization analysis

We undertook a two-sample Mendelian randomization approach²⁷ to estimate the causal effect of TB-BMD on any-type of fracture in the Europeans samples. In short, we constructed a score based on the independent genetic variants from the TB-BMD meta-analysis (European set and excluding secondary signals), whenever the selected variant was not present in the fracture meta-analysis, the second variant with the lowest p-value in the locus ($P < 5 \times 10^{-8}$) and $r^2 > 0.8$ was used as proxy. Thereafter, estimates derived from the TB-BMD summary statistics were pooled using methods similar to inverse-variance weighted fixed meta-analysis using the meta R-package (**Web Resources**).

Search for biological and functional knowledge of the identified association regions

For all those SNPs outside a 500Kb window from previously known bone associated SNPs we did a literature search in PubMed and Web of Science to evaluate if nearby genes (within 500Kb) were known to play a role in bone metabolism. Also, we determined if the annotated genes underlie any human Mendelian disorder with a skeletal manifestation, had knockout mouse models with a skeletal phenotype or were annotated to pathways critical to bone metabolism. Genomic annotation for all SNPs was made based on UCSC hg19.

DEPICT analyses

We used DEPICT²⁸, a recently developed tool to prioritize genes at the associated regions, define possible pathways by enrichment testing, and identify tissue and cell types in which genes from loci associated with TB-BMD. The methodology first selects all lead SNPs below a certain threshold with respect to a target P-value. We tested both the complete set of GWS

SNPs and the subset of those mapping only to loci not previously reported. Enriched gene-set were group based on the degree of gene overlap into ‘meta gene-sets’ as proposed earlier²⁹, and their correlation visualized using Cytoscape 3.4 (**Web Resources**).

Functional annotation to microRNA binding sites

We used the PolymiRTS²⁹, miRdSNP³⁰, and microSNiPer³¹ databases to obtain a list of variants located in predicted microRNA binding sites on the 3’UTRs of genes, as described in detail elsewhere³². In summary, index SNPs (most associated variant) of the GWS loci were submitted to SNAP (**Web Resources**) to retrieve their high LD proxy SNPs (with $r^2 > 0.8$, limit distance 500 kb, and CEU panel) in the 1000 genomes project. The resulting list of SNPs was annotated to the list of microRNA binding site variants obtained from the above mentioned publicly available databases.

Functional enrichment analysis of trait-associated variants

GWAS Analysis of Regulatory or Functional Information Enrichment with LD correction (GARFIELD)³³ was used to characterize the putative functional contribution of TB-BMD associated variants mapping to non-coding regions. GARFIELD employs a non-parametric analysis to calculate fold enrichment values for regulatory marks, at given significance thresholds and then tests them for significance via permutation testing while accounting for LD, MAF and local gene density³³. We used data regarding DNase I hypersensitive sites, transcription factor binding sites, histone modifications and chromatin states (ENCODE and Roadmap Epigenomics) from 424 cell types and tissues to capture and characterize possible cell-type-specific patterns of enrichment, as provided in the GARFIELD software (**Web**

Resources). Fold enrichment statistics were tested at the four different significance thresholds (i.e., 1×10^{-8} , 1×10^{-7} , 1×10^{-6} and 1×10^{-5}). Multiple-testing correction was performed on the effective number of annotations used, using the default P-value threshold of 1×10^{-4} .

Knockout animal models and gene expression in bone cells

Animal models survey

We surveyed databases from *The International Mouse Phenotyping Consortium*³⁴ together with *The International Knockout Mouse Consortium*³⁵ to identify knockout models of candidate genes resulting in skeletal phenotypes. Furthermore we mined data from *The Origins of Bone and Cartilage Disease* (OBCD) project³⁶, specialized in murine skeletal phenotypes including Digital X-ray microradiography on femurs and tail vertebrae, Micro-CT analysis, femur three-point bend test load–displacement curves and tail vertebrae compression testing from knockout mice and wild-type controls at 16 weeks of age.

Gene expression in murine bone cells

Gene expression profiles of candidate genes were examined in primary mouse osteoblasts undergoing differentiation and bone marrow derived osteoclasts. To study murine osteoblasts, pre-osteoblast-like cells were obtained from neonatal calvaria collected from C57BL/6J. Next Generation RNA sequencing using an Illumina HiSeq 2000 was used to evaluate the transcriptome every two days from day 2 to 18 days post osteoblast differentiation⁷. Expression of genes in murine osteoclasts was determined using publicly available data obtained using Next-Gen RNA-sequencing applied to bone marrow derived osteoclasts obtained from 6-8 week old C57BL/6 mice³⁷.

Gene expression in human bone cells

Gene expression profiles of candidate genes were examined in human bone marrow derived mesenchymal stem cells differentiated into osteoblast. Total RNA (n=3) was isolated at day 0 (MSCs) and day 4 of osteoblast differentiation³⁸. Also, RNA was isolated during osteoclast differentiation. Peripheral blood mononuclear cells derived from buffy coats (Sanquin, Amsterdam, the Netherlands) were seeded in 96-well plates (5×10^5 cells per well) as previously described³⁹. Total RNA (n=3) was isolated using Trizol at day 0 (PBMCs) and at day 7 of osteoclast differentiation. Illumina HumanHT-12 v3 BeadChip human whole-genome expression arrays were used for expression profiling. The quality of isolated RNA was assessed on a 2100 Bioanalyzer (Agilent Technologies). Data were analyzed as described in detail previously³⁸. Genes were designated as being expressed when at least one probe coding for the gene was significantly present in at least 2 of the 3 biological replicates.

Results

TB-BMD GWAS meta-analyses

Analyses including all age-strata

Our meta-analysis of TB-BMD GWAS summary statistics (N=66,628) identified variants in 76 independent loci associated with TB-BMD at a genome-wide significant (GWS, $P \leq 5 \times 10^{-8}$) level (**Figure 1, Table S4**). Overall, there was no evidence of a strong inflation (genomic inflation factor (λ) of 1.08, **Figure S1**). Yet, inflation was observed in the range of common variants ($0.2 > \text{MAF} < 0.5$, $\lambda = 1.19$) due to polygenicity (LD score regression intercept = 1.007). In our results, one of the signals mapping to *LDLRAD3* was driven entirely by individuals of African

background (MAF=0.043 in YRI panel) since the two associated variants are monomorphic in all other populations. The low allele frequency of this variant in our study (MAF= 0.025) and our limited statistical power (N=6,748) in non-European samples warrants independent replication efforts to exclude the possibility of a false-positive association.

In addition, a meta-analysis comprising 56,284 individuals of European ancestry (~84% of the study population) identified variants in two additional GWS loci (**Figures S1-S2, Table S5**). Association signals mapping to these loci were close to the GWS threshold in the overall meta-analysis ($P=1\times 10^{-7}$) and showed no evidence of heterogeneity ($P_{\text{het}}>0.1$). One of them, in 12q24.21 (*MED13L*), has not been previously associated with bone parameters (**Table 1, Figure S3**), while the other in 21q22.13 (*CLDN14*), is not fully independent from the previously reported hip-BMD association signal¹³ (**Table S5**).

Of the 78 identified loci, variants in 35 (45%) were not located within 500 kb of known association signals nor in regions of extended LD with them (**Table 1, Figure S4**). Index SNPs at these 35 loci were, in general, common non-coding variants. Twenty-two of these, are located in close proximity to genes likely to influence bone metabolism as shown by previous functional studies (**Table 1, Figure S3**), including *CSF1* ([MIM 120420] important for osteoclast differentiation⁴⁰) and *SMAD3* ([MIM 603109] a critical component of the TGF-beta signaling pathway⁴¹). Across these 35 signals, 31 of the index SNPs were nominally associated ($P<0.05$) with either lumbar spine or femoral neck BMD in the same direction as in the previously published GEFOS GWAS meta-analysis⁷ (**Table 1**). This comparison was not possible for the rs113964474 variant, because it was not available in the GEFOS study. Moreover, we found directionally-concordant effect estimates ($P < 0.05$) for 73 of the 78 index SNPs of known bone

association signals (**Table S3**). The markers which failed to replicate in our study were either previously associated with lumbar spine BMD but not femoral neck BMD (rs3905706 [*MPP7*, 10p12.1] and rs1878526 [*INSIG2*, 2q14.2]), associated specifically with the hip trochanter and intertrochanteric subregions (rs1949542 [*RP11-384F7.1*, 3q13.32]), or associated with BMD only in women (rs7017914 [*XKR9*, 8q13.3]) or only in children (rs754388 [*RIN3*, 14q32.12]).

Age-dependent effects

Meta-analyses across age strata resulted in the identification of variants mapping to 2 additional loci that were not detected in the overall meta-analysis (**Figure S5; Table S6**). In children (age group 0-15 years), the previously known 14q32.12 locus⁸, harboring *RIN3* (rs72699866, $P=1\times 10^{-8}$); and in the middle-aged (age group 45-60 years), a signal in the 19q12 locus mapping in the vicinity of *TSHZ3* (rs6510186, $P=3.1\times 10^{-8}$) were identified. The rs72699866 variant leading the *RIN3* signal in the youngest age stratum showed no evidence of association ($P=0.16$) and high heterogeneity ($P_{\text{het}}=6.6\times 10^{-5}$) in the overall meta-analysis. In fact, the effect of rs72699866 decreased significantly with age ($P_{\text{trend}}=1.69\times 10^{-9}$) (**Figure S6**) and showed a significant difference between the two extreme groups, i.e. children vs elderly ($\beta_{0-15}=0.099$ [0.066, 0.134]; $\beta_{>60}=-0.035$ [-0.060, -0.010]; $P_{\text{diff}}=4.32\times 10^{-10}$). In contrast, the rs6510186 variant [19q12] showed nominal evidence of association and heterogeneity in the overall meta-analysis ($P=0.02$; $P_{\text{het}}=0.03$). Nevertheless, no clear pattern of age-dependency was identified ($P=0.2$) for this SNP (**Figure S6**).

We also applied meta-regression analysis and found that variants mapping to 42 different loci showed nominally significant age dependent effect ($P<0.05$) (**Table S7, Figure S7**). In summary,

27 (64%) of the loci showed stronger effects in the older age groups. Of these, variants in the 6q25.1 (*ESR1*) and 13q14.11 (*RANKL*) loci remained significant after multiple-testing correction ($P < 3.4 \times 10^{-5}$) (**Figure 2**); while variants in 6p21.1 (*RUNX2*, rs148460475), 15q21.2 (*CYP19A1*, rs2414098), 17q21.31 (*MEOX1*, rs74835612) and 11p15.1 (*SOX6*, rs11822790) were only suggestive at $P < 1 \times 10^{-3}$.

Conditional association analyses

The step-wise conditional approach included studies comprising only individuals of European ancestry, as the method used relies on appropriate representability of the LD reference. Of the 76 GWS loci identified in the overall analysis, variants in 57 (19 previously unreported) loci were also GWS in the European-only analysis (**Figure S2**), likely a consequence of the lower power in this subgroup. We identified 81 SNPs independently associated with TB-BMD mapping to 58 different loci (one European-specific), 18 of which depicted multiple distinct signals attaining GWS (**Table S8**). These independent variants together explained 10.2% of TB-BMD variance. This proportion is slightly higher than the 7.4% TB-BMD variance explained by the 78 known variants associated with bone traits. Moreover, we identified independent signals in 13 of the 78 known bone loci after conditional analyses. (**Figure S2; Table S8**).

Shared Genetic architecture of TB-BMD, fracture and other traits

SNP-heritability of TB-BMD in the European samples was estimated to be 0.259 (SE 0.017). TB-BMD was highly genetically correlated with BMD measured at other skeletal sites ($\rho > 0.9$). Among the non-BMD traits, all-type of fracture showed the highest correlation [$\rho = -0.61$ ($P = 1.6 \times 10^{-27}$)]. The MR approach indicated a strong causal relation where per 1 standard

deviation decrease in genetically determined TB-BMD there is 56% increase in the risk of fracture (Odds ratio 1.56 [1.50-1.62]). Other anthropometric, metabolic and disease traits showed significant (yet weak) correlation with TB-BMD (**Table S9, Figure 3**). In contrast, other established risk factors for osteoporosis such as menopause or age of menarche showed no significant genetic correlation with TB-BMD.

Biological and functional knowledge of the genes in BMD-associated loci

Loci not previously reported and their potential role in bone metabolism are summarized in **Table 1**. Several loci harbor genes implicated directly in bone metabolism (*SLC8A1* [MIM 182305], *PLCL1* [MIM 600597], *ADAMTS5* [MIM 605007]), affecting osteoblast or osteoclast differentiation and activity (*CSF1* [MIM 120420], *DUSP5* [MIM 603069], *SMAD3* [MIM 603109], *SMAD9* [MIM 603295], *CD44* [MIM 107269]), participating in Wnt signaling (*FZD7* [MIM 603410], *TCF7L1* [MIM 604652]), or regulating processes such as manganese or calcium absorption (*GCKR* [MIM 600842], *DGKD* [MIM 601826], *SLC30A10* [MIM 611146]) among others⁴⁰⁻⁶¹; while genes in at least 14 loci exert a potential novel role in bone biology. Rodent knockout models of several genes in the implicated loci, show an altered skeletal phenotype (e.g., osteopetrosis [*Csf1*⁴⁰], increased bone resorption [*Aqp1*⁵⁰, *Cyp19a1*⁵⁷, *Cd44*⁵³], impaired skeletogenesis [*Apc*⁴⁹, *Runx1*⁶⁰, *Smad3*⁴¹], deformities in the axial skeleton [*Btg1*⁶², *Atpaf2*⁶³]). Whereas an effect on bone can be inferred for genes in other associated loci, for example, *CYP19A1* [MIM 107910] in 15q21.2 is an estrogen synthesis gene, being estrogen a key compound for bone maturation and maintenance, and *ZKSCAN5* [MIM 611272] in 7q22.1 is associated with circulating dehydroepiandrosterone sulphate (DHEAS) levels⁵¹. DHEAS levels are positively correlated with BMD in adults and post-menopausal women⁶⁴. Across these loci,

not previously reported as associated with BMD variation, we identified six exonic variants associated with TB-BMD, three of which were nonsynonymous variants all cataloged as benign both by SIFT and polyphen2. We also identified 53 GWS coding variants in known loci, of which 33 are non-synonymous (**Table S10**). Only a low-frequency variant in *LRP5* [MIM 603506], rs4988321/A (11:68174189, MAF=0.04), has a clinical annotation, constituting a homozygous G-to-A transition variant identified in a person with osteoporosis-pseudoglioma syndrome (OPPG [MIM 259770])⁶⁵.

DEPICT analyses

Based on the overall meta-analysis, 53 genes were prioritized (FDR<0.05), 15 of them mapping to loci not previously described (**Table S11**). Cells and tissues from the musculoskeletal system presented the largest enrichment of gene expression within the associated loci (**Figure 4**). These genes were overrepresented in 182 pathways clustered in 25 ‘meta gene-sets’ (**Table S12**). The large majority of the clusters are involved in musculoskeletal development and bone homeostasis (**Figure 4**). The most significant of these implicated the regulation of cell growth, and the TGFB signaling pathway and its mediating SMAD proteins.

Restricting the DEPICT analysis to the subset of not previously reported associated regions resulted in significant enrichment of genes expressed in the musculoskeletal and immunological systems (**Figure S8**). Genes mapping to these loci were overrepresented in the SMAD binding pathway and TGFBR2 PPI (protein-protein interaction) subnetwork (FDR<0.05).

Functional annotation to microRNA binding sites

We then assessed if the index SNPs of the 80 GWS loci detected in the main and subsequent GWAS (or their proxies in strong LD; $r^2 > 0.8$) were located in predicted microRNA binding sites within the genes' 3'UTRs and thus, were expected to disrupt the regulation of gene expression (**Table S13**). The index SNP within the 3'UTR of *ZKSCAN5* (mapping to a locus not previously identified), rs34670419 (MAF=0.04), is predicted to create a binding site for miR-382-3p, a microRNA which is expressed in osteocytes and has been recently shown to be involved in osteogenic differentiation⁶⁶. In addition, eight proxy SNPs (mapping to *PSMD13*, *ABCF2*, *GALNT3*, *PKDCC*, *REEP5*, *PPP6R3*, *AAGAB* and *TOM1L2*) are predicted to influence the binding of microRNAs to transcripts of their host gene.

Functional enrichment analysis of trait-associated variants

As typically found in GWAS, the great majority of identified associations emerged from non-coding common variants and hold no direct annotation to molecular mechanisms. To assess if there is relative enrichment of regulatory genomic marks underlying the associated variants in a cell-specific context, we used GARFIELD³³. We found relative ubiquitous enrichment for TB-BMD variants (Empirical $P < 2.4 \times 10^{-4}$) in DNase I hypersensitive sites across the different cell types (**Figure S9**). Further, we found higher levels of fold-enrichment for enhancers (median 3.6, range [2.7, 4.4]) and promoters (median 3.2, range [2.9, 3.5]) than for transcribed regions (median 1.8, range [1.5, 2.2]).

Gene expression in bone cells and knockout animal models

From the 53 genes prioritized by DEPICT only 49 had a mouse orthologue (**Table S14**). From these genes, only *Mepe* (osteocyte-specific) and *Foxl1* were not expressed in murine osteoblast

or osteoclast. Moreover, 61% of the prioritized genes were expressed in human cells *in vitro* during osteoblast or osteoclast differentiation (**Table S14**). *AQP1* was the only prioritized gene mapping to a locus not previously reported showing no expression in the human bone cells differentiation experiments.

Knockout models were widely available in at least one of the different databases assessed. Nevertheless in-depth bone phenotyping performed under the OBCD project was only available for four knockout models (**Table S15**). Two of these, *DUSP5* and *CD300LG* showed no significant bone phenotype. The *TCF7L1* knockout model only showed lower cortical diameter in the femur without other clear bone phenotype. Nevertheless, *TCF7L1* was shown to be expressed during osteoblastogenesis. Conversely, homozygous knockout for *CREB3L1* showed a clear bone phenotype consisting of low BMC both at the vertebrae and femur together with a strong trabecular and cortical phenotype affecting bone strength (**Figure S10**). *CREB3L1* maps to 11p11.2, a previously identified BMD locus⁶ harboring *ARHGAP1* and *LRP4* as candidates to underlie the GWAS signal in a region of extended LD.

Discussion

This meta-analysis of TB-BMD comprising up to 66,000 individuals identified variants in 36 loci not previously reported and replicated at GWS level several association signals identified by GWAS of diverse bone phenotypes. Bioinformatics analyses suggest enrichment of these 36 loci for genes expressed in the musculoskeletal system, and solidly represented in the SMAD binding pathway and the TGFBR2 PPI subnetwork. We also demonstrate that for variants in few loci the size of the effect is age dependent; variants in two loci (*RIN3* and *TSHZ3*) were

identified only by the age-stratified analyses despite less power (smaller sample size); while for variants in two other loci (*ESR1* and *RANKL*) there was significant evidence of age heterogeneity derived from a meta-regression of the genetic effects with age. Our results strengthen the evidence that genetic variants influence BMD from a young age and support the value of peak bone mass as an important determinant of bone health later in life.

Traditionally, DXA-BMD measurements performed at sites of high fracture risk (i.e., femoral neck, lumbar spine and forearm) have been used in genetic epidemiological investigations of bone health in adults. Instead, we have used BMD measurements derived from total body scans. Not only do we show a high overlap of association signals with previous GWAS of different bone traits, including DXA, pQCT and ultrasound measurements, but we have also identified unreported loci. Five known associations failed to replicate in our studies, even though we cannot discard these associations constitute false-positives, these results might also indicate that variants whose effect is highly specific to skeletal sites, skeletal properties, sex or age groups cannot be detected in our TB-BMD meta-analysis. It is plausible that more variants of this type exist and will be discovered as site-specific BMD meta-analyses are performed in increasingly powered settings. Furthermore, the genetic correlation of TB-BMD with BMD measured at other sites was close to one. Whilst, we found that a decrease of one standard deviation in the genetically determined TB-BMD resulted in at least 50% higher odds of suffering a fracture. Significant genetic correlations with other traits (i.e., BMI, IGF1 and ulcerative colitis) reflect the systemic context of skeletal biology and merit further study by future efforts to elucidate the underlying mechanisms.

505 Genes in the associated loci were highly expressed in the musculoskeletal system and
506 overrepresented in gene-sets related to bone development. The prioritized gene *CREB3L1* [MIM
507 616215] in 11p11.2 observed a clear bone phenotype in our mouse knockout model, which
508 corroborates the findings of previous work showing substantial rescue of *CREB3L1* deficiency
509 with bisphosphonates and its critical role for bone formation⁶⁷. This locus characterized by
510 extended LD, also harbors *LRP4* [MIM 604270] whose knockout model presents with increased
511 trabecular and cortical bone mass⁶⁸. This is in line with our conditional analysis identifying
512 multiple independent signals in the region making it likely that both genes are influencing bone
513 biology. Altogether, we demonstrated that TB-BMD offers a powerful alternative to identify
514 genetic variants associated with bone metabolism.

515 Variants mapping to 14q32 harboring *RIN3* [MIM 610223] were only associated at a GWS level
516 in children (i.e., <15 years), and were only nominally significant in the elderly group (i.e., >60
517 years). This age-related heterogeneity may explain why this locus has not been detected in
518 BMD meta-analyses in adults, although being identified in relation to pediatric BMD⁸ and
519 Paget's disease (PDB [602080]) GWAS⁶⁹. In addition, another signal mapping to 19q12
520 harboring *TSHZ3* [MIM 614119] was significant in adults aged 45-60 years but not in other age
521 groups analyzed or in previous studies, alluding to a false-positive association, thus replication
522 of this finding is necessary.

523 Our analyses revealed variants in the 6q25.1 (*ESR1*) and 13q14.11 (*RANKL*) loci demonstrating
524 the most compelling evidence for age-modulation effects. The 6q25.1 locus harboring *ESR1*
525 [MIM 133430], an important genetic factor in normal BMD variability, was not associated with
526 BMD in children below 15 years of age, where the largest cohorts (i.e., Avon Longitudinal Study

527 of Parents and Children (ALSPAC) and the Generation R Study) comprise predominantly pre-
528 pubertal children. As levels of estradiol before puberty are low⁷⁰, a negligible effect of *ESR1*
529 variants on BMD is expected. Likewise, in mouse models the expression of *RANKL* [MIM
530 602642] in bone is markedly increased with advancing age from young to adult and related to
531 bone loss⁷¹. Accordingly, variants influencing *RANKL* expression show a larger effect later in life.
532 In general, a substantial heterogeneity of the genetic effects in the overall meta-analysis was
533 explained by age, nevertheless, the inclusion of larger sample sizes (avoiding age exclusion
534 criteria and incrementing statistical power) leveled off the loss of power due to the
535 heterogeneity of the genetic effects.

536 In brief, variants with evidence of age-specific effects were exceptional in our study. These
537 results might reflect a lack of statistical power as only SNPs showing suggestive evidence
538 ($P < 5 \times 10^{-6}$) of association with TB-BMD in the overall meta-analysis were tested for age-specific
539 effects. This selection criteria aimed to include SNPs whose heterogeneity might have
540 hampered their statistical significance in the overall meta-analysis, and at the same time
541 maximize the power to discover variants with real age-dependent effects. Alternatively, these
542 results indicate that most of the genetic variants identified so far, by us and others, influence
543 BMD from early ages onwards, and their effect persist throughout the life course. However,
544 variants in 27 of the 42 loci (64%) showing nominal evidence for age dependent effects had
545 larger effects in the older groups. Nonetheless, this requires careful interpretation given the
546 uneven sample sizes between the age groups and the criteria to select markers for the meta-
547 regression based on significance in the overall meta-analysis. Collectively, this argues in favor of

enlarging studies focused on younger populations –where the statistical power is still restricted – to discover additional genetic variants influencing BMD.

Our study has some limitations. A key disadvantage of our design is that we group the data based on age spans rather than life stages. Crucial information for this assesment, such as puberty onset in children and adolescents or menopausal status in the adults, was not available across the majority of the cohorts. Other strategies like using shorter age spans will resulted in even less statistical power of the discovery setting. Similarly, despite the large sample size of our study, we identified very few variants in the low-frequency spectrum (MAF <5%) indicating that comprehensive surveys of rare variation influencing BMD still require even larger sample sizes, on top of better resources for imputation of the rarer variants, possibly needing population-specific references. Such strategies will be key to explain a larger fraction of the genetic variability of BMD phenotypes, as illustrated for other traits such as height or BMI⁷². Moreover, the identified SNPs are in their vast majority, non-coding variants, raising the possibility that the causal genes are different from the candidate genes we have prioritized based on the current biological knowledge and bioinformatic prediction tools. Additional functional studies are required to determine the potential role of the genes in the identified loci.

In conclusion, we performed a genome-wide survey for association with DXA derived TB-BMD, combining data from five age groups including children and older individuals. In contrast to previous large-scale meta-analyses^{6;7}, we used DXA derived TB-BMD rather than measurements on specific skeletal sites prone to fracture to identify genetic factors influencing BMD variation. We demonstrate that TB-BMD is a valid phenotype for this purpose, as we replicated more than

90% of the previously reported signals. Most importantly, we identify variants in 36 loci associated with TB-BMD not previously reported by previous GWAS of bone phenotypes. Our results show steadiness in the magnitude of the genetic effects on BMD for most of the BMD-associated variants. While the contrasting skeletal physiology across different age periods is well established (i.e. endochondral ossification, linear growth, modelling, remodeling, etc.), peak bone mass acquisition remains the major determinant of variability at any age. These findings strongly support the importance of the bone accrual process in the definition of BMD status and fracture susceptibility throughout the life course.

Accession Numbers

GWAS Summary data for the main and age-strata meta-analyses together with the corresponding regional plots of GWS signals have been deposited in the GEFOS website (**Web Resources**). Gene expression data presented in this paper can be retrieved from the Gene Expression Omnibus (GEO) as follows: Murine osteoclasts (GSM1873361) and osteoblasts (GSE54461); human osteoblast differentiation (GSE54461).

Supplemental Data

Supplemental data include a full list of acknowledgements, cohort short descriptions, 15 tables and 10 figures.

Acknowledgements

The authors would like to thank the many colleagues who contributed to collection and phenotypic characterization of the clinical samples, as well as genotyping and analysis of the

590 GWAS data. Part of this work was conducted using the UK Biobank resource.

591 **Conflict of interests**

592 Psaty serves on the DSMB of a clinical trial for the manufacturer (Zoll LifeCor) and on the

593 Steering Committee of the Yale Open Data Access Project funded by Johnson & Johnson.

594

595 **Web Resources**

596 GARFIELD, <http://www.ebi.ac.uk/birney-srv/GARFIELD/>GEFOS, <http://www.gefos.org/>

597 LDhub, <http://ldsc.broadinstitute.org/>

598 Meta R-package, <https://github.com/guido-s/meta>

599 OBCD, <http://www.boneandcartilage.com/>

600 OMIM, <http://www.omim.org/>

601 SNAP, <http://archive.broadinstitute.org/mpg/snap/>

References

1. Johnell, O., Kanis, J.A., Oden, A., Johansson, H., De Laet, C., Delmas, P., Eisman, J.A., Fujiwara, S., Kroger, H., Mellstrom, D., et al. (2005). Predictive value of BMD for hip and other fractures. *J Bone Miner Res* 20, 1185-1194.
2. Farr, J.N., and Khosla, S. (2015). Skeletal changes through the lifespan--from growth to senescence. *Nat Rev Endocrinol* 11, 513-521.
3. Hendrickx, G., Boudin, E., and Van Hul, W. (2015). A look behind the scenes: the risk and pathogenesis of primary osteoporosis. *Nat Rev Rheumatol* 11, 462-474.
4. International Society for Clinical Densitometry. (2015). Official ISCD Positions – Adult & Pediatric individuals, <http://www.iscd.org/official-positions/>
5. Tenne, M., McGuigan, F., Besjakov, J., Gerdhem, P., and Akesson, K. (2013). Degenerative changes at the lumbar spine--implications for bone mineral density measurement in elderly women. *Osteoporos Int* 24, 1419-1428.
6. Estrada, K., Styrkarsdottir, U., Evangelou, E., Hsu, Y.H., Duncan, E.L., Ntzani, E.E., Oei, L., Albagha, O.M., Amin, N., Kemp, J.P., et al. (2012). Genome-wide meta-analysis identifies 56 bone mineral density loci and reveals 14 loci associated with risk of fracture. *Nat Genet* 44, 491-501.
7. Zheng, H.F., Forgetta, V., Hsu, Y.H., Estrada, K., Rosello-Diez, A., Leo, P.J., Dahia, C.L., Park-Min, K.H., Tobias, J.H., Kooperberg, C., et al. (2015). Whole-genome sequencing identifies EN1 as a determinant of bone density and fracture. *Nature* 526, 112-117.
8. Kemp, J.P., Medina-Gomez, C., Estrada, K., St Pourcain, B., Heppe, D.H., Warrington, N.M., Oei, L., Ring, S.M., Kruithof, C.J., Timpson, N.J., et al. (2014). Phenotypic dissection of bone mineral density reveals skeletal site specificity and facilitates the identification of novel loci in the genetic regulation of bone mass attainment. *PLoS Genet* 10, e1004423.
9. Paternoster, L., Lorentzon, M., Lehtimäki, T., Eriksson, J., Kahonen, M., Raitakari, O., Laaksonen, M., Sievanen, H., Viikari, J., Lyytikäinen, L.P., et al. (2013). Genetic determinants of trabecular and cortical volumetric bone mineral densities and bone microstructure. *PLoS Genet* 9, e1003247.
10. Moayyeri, A., Hsu, Y.H., Karasik, D., Estrada, K., Xiao, S.M., Nielson, C., Srikanth, P., Giroux, S., Wilson, S.G., Zheng, H.F., et al. (2014). Genetic determinants of heel bone properties: genome-wide association meta-analysis and replication in the GEFOS/GENOMOS consortium. *Hum Mol Genet* 23, 3054-3068.

11. Medina-Gomez, C., Kemp, J.P., Estrada, K., Eriksson, J., Liu, J., Reppe, S., Evans, D.M., Høpfe, D.H., Vandenput, L., Herrera, L., et al. (2012). Meta-analysis of genome-wide scans for total body BMD in children and adults reveals allelic heterogeneity and age-specific effects at the WNT16 locus. *PLoS Genet* 8, e1002718.
12. Yang, T.L., Guo, Y., Liu, Y.J., Shen, H., Liu, Y.Z., Lei, S.F., Li, J., Tian, Q., and Deng, H.W. (2012). Genetic variants in the SOX6 gene are associated with bone mineral density in both Caucasian and Chinese populations. *Osteoporos Int* 23, 781-787.
13. Zhang, L., Choi, H.J., Estrada, K., Leo, P.J., Li, J., Pei, Y.F., Zhang, Y., Lin, Y., Shen, H., Liu, Y.Z., et al. (2014). Multistage genome-wide association meta-analyses identified two new loci for bone mineral density. *Hum Mol Genet* 23, 1923-1933.
14. Pei, Y.F., Xie, Z.G., Wang, X.Y., Hu, W.Z., Li, L.B., Ran, S., Lin, Y., Hai, R., Shen, H., Tian, Q., et al. (2016). Association of 3q13.32 variants with hip trochanter and intertrochanter bone mineral density identified by a genome-wide association study. *Osteoporos Int*.
15. Styrkarsdottir, U., Thorleifsson, G., Eiriksdottir, B., Gudjonsson, S.A., Ingvarsson, T., Center, J.R., Nguyen, T.V., Eisman, J.A., Christiansen, C., Thorsteinsdottir, U., et al. (2016). Two Rare Mutations in the COL1A2 Gene Associate With Low Bone Mineral Density and Fractures in Iceland. *J Bone Miner Res* 31, 173-179.
16. Koller, D.L., Zheng, H.F., Karasik, D., Yerges-Armstrong, L., Liu, C.T., McGuigan, F., Kemp, J.P., Giroux, S., Lai, D., Edenberg, H.J., et al. (2013). Meta-analysis of genome-wide studies identifies WNT16 and ESR1 SNPs associated with bone mineral density in premenopausal women. *J Bone Miner Res* 28, 547-558.
17. Nielson, C.M., Liu, C.T., Smith, A.V., Ackert-Bicknell, C.L., Reppe, S., Jakobsdottir, J., Wassel, C., Register, T.C., Oei, L., Alonso, N., et al. (2016). Novel Genetic Variants Associated With Increased Vertebral Volumetric BMD, Reduced Vertebral Fracture Risk, and Increased Expression of SLC1A3 and EPHB2. *J Bone Miner Res* 31, 2085-2097.
18. Styrkarsdottir, U., Thorleifsson, G., Gudjonsson, S.A., Sigurdsson, A., Center, J.R., Lee, S.H., Nguyen, T.V., Kwok, T.C., Lee, J.S., Ho, S.C., et al. (2016). Sequence variants in the PTCH1 gene associate with spine bone mineral density and osteoporotic fractures. *Nat Commun* 7, 10129.
19. Winkler, T.W., Day, F.R., Croteau-Chonka, D.C., Wood, A.R., Locke, A.E., Magi, R., Ferreira, T., Fall, T., Graff, M., Justice, A.E., et al. (2014). Quality control and conduct of genome-wide association meta-analyses. *Nat Protoc* 9, 1192-1212.

20. Simino, J., Shi, G., Bis, J.C., Chasman, D.I., Ehret, G.B., Gu, X., Guo, X., Hwang, S.J., Sijbrands, E., Smith, A.V., et al. (2014). Gene-age interactions in blood pressure regulation: a large-scale investigation with the CHARGE, Global BPgen, and ICBP Consortia. *Am J Hum Genet* 95, 24-38.
21. Viechtbauer, W. (2010). Conducting Meta-Analyses in R with the metafor Package. *J Stat Softw* 36, 1-48.
22. Winkler, T.W., Kutalik, Z., Gorski, M., Lottaz, C., Kronenberg, F., and Heid, I.M. (2015). EasyStrata: evaluation and visualization of stratified genome-wide association meta-analysis data. *Bioinformatics* 31, 259-261.
23. Yang, J., Ferreira, T., Morris, A.P., Medland, S.E., Madden, P.A.F., Heath, A.C., Martin, N.G., Montgomery, G.W., Weedon, M.N., Loos, R.J., et al. (2012). Conditional and joint multiple-SNP analysis of GWAS summary statistics identifies additional variants influencing complex traits. *Nature Genetics* 44, 369-U170.
24. Bulik-Sullivan, B.K., Loh, P.R., Finucane, H.K., Ripke, S., Yang, J., Schizophrenia Working Group of the Psychiatric Genomics, C., Patterson, N., Daly, M.J., Price, A.L., and Neale, B.M. (2015). LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet* 47, 291-295.
25. Bulik-Sullivan, B., Finucane, H.K., Anttila, V., Gusev, A., Day, F.R., Loh, P.R., ReproGen, C., Psychiatric Genomics, C., Genetic Consortium for Anorexia Nervosa of the Wellcome Trust Case Control, C., Duncan, L., et al. (2015). An atlas of genetic correlations across human diseases and traits. *Nat Genet* 47, 1236-1241.
26. Zheng, J., Erzurumluoglu, A.M., Elsworth, B.L., Kemp, J.P., Howe, L., Haycock, P.C., Hemani, G., Tansey, K., Laurin, C., Early, G., et al. (2017). LD Hub: a centralized database and web interface to perform LD score regression that maximizes the potential of summary level GWAS data for SNP heritability and genetic correlation analysis. *Bioinformatics* 33, 272-279.
27. Burgess, S., Butterworth, A., and Thompson, S.G. (2013). Mendelian randomization analysis with multiple genetic variants using summarized data. *Genet Epidemiol* 37, 658-665.
28. Pers, T.H., Karjalainen, J.M., Chan, Y., Westra, H.J., Wood, A.R., Yang, J., Lui, J.C., Vedantam, S., Gustafsson, S., Esko, T., et al. (2015). Biological interpretation of genome-wide association studies using predicted gene functions. *Nature Communications* 6.
29. Bhattacharya, A., Ziebarth, J.D., and Cui, Y. (2014). PolymiRTS Database 3.0: linking polymorphisms in microRNAs and their target sites with human diseases and biological pathways. *Nucleic Acids Res* 42, D86-91.

30. Gong, J., Tong, Y., Zhang, H.M., Wang, K., Hu, T., Shan, G., Sun, J., and Guo, A.Y. (2012). Genome-wide identification of SNPs in microRNA genes and the SNP effects on microRNA target binding and biogenesis. *Hum Mutat* 33, 254-263.
31. Barenboim, M., Zoltick, B.J., Guo, Y., and Weinberger, D.R. (2010). MicroSNIPer: a web tool for prediction of SNP effects on putative microRNA targets. *Hum Mutat* 31, 1223-1232.
32. Ghanbari, M., Franco, O.H., de Looper, H.W., Hofman, A., Erkeland, S.J., and Dehghan, A. (2015). Genetic Variations in MicroRNA-Binding Sites Affect MicroRNA-Mediated Regulation of Several Genes Associated With Cardio-metabolic Phenotypes. *Circ Cardiovasc Genet* 8, 473-486.
33. Iotchkova, V., Ritchie, G.R.S., Geijs, M., Morganella, S., Min, J.L., Walter, K., Timpson, N.J., Dunham, I., Birney, E., and Soranzo, N. (2016). GARFIELD - GWAS Analysis of Regulatory or Functional Information Enrichment with LD correction. *bioRxiv*.
34. Dickinson, M.E., Flenniken, A.M., Ji, X., Teboul, L., Wong, M.D., White, J.K., Meehan, T.F., Weninger, W.J., Westerberg, H., Adissu, H., et al. (2016). High-throughput discovery of novel developmental phenotypes. *Nature* 537, 508-514.
35. Hrabe de Angelis, M., Nicholson, G., Selloum, M., White, J.K., Morgan, H., Ramirez-Solis, R., Sorg, T., Wells, S., Fuchs, H., Fray, M., et al. (2015). Analysis of mammalian gene function through broad-based phenotypic screens across a consortium of mouse clinics. *Nat Genet* 47, 969-978.
36. Freudenthal, B., Logan, J., Sanger Institute Mouse, P., Croucher, P.I., Williams, G.R., and Bassett, J.H. (2016). Rapid phenotyping of knockout mice to identify genetic determinants of bone strength. *J Endocrinol* 231, R31-46.
37. Kim, K., Punj, V., Kim, J.M., Lee, S., Ulmer, T.S., Lu, W., Rice, J.C., and An, W. (2016). MMP-9 facilitates selective proteolysis of the histone H3 tail at genes necessary for proficient osteoclastogenesis. *Genes Dev* 30, 208-219.
38. van de Peppel, J., Strini, T., Tilburg, J., Westerhoff, H., van Wijnen, A.J., and van Leeuwen, J.P. (2017). Identification of Three Early Phases of Cell-Fate Determination during Osteogenic and Adipogenic Differentiation by Transcription Factor Dynamics. *Stem Cell Reports* 8, 947-960.
39. Koek, W.N.H., van der Eerden, B.C.J., Alves, R., van Driel, M., Schreuders-Koedam, M., Zillikens, M.C., and van Leeuwen, J. (2017). Osteoclastogenic capacity of peripheral blood mononuclear cells is not different between women with and without osteoporosis. *Bone* 95, 108-114.
40. Dobbins, D.E., Sood, R., Hashiramoto, A., Hansen, C.T., Wilder, R.L., and Remmers, E.F. (2002). Mutation of macrophage colony stimulating factor (Csf1) causes osteopetrosis in the tl rat. *Biochem Biophys Res Commun* 294, 1114-1120.

41. Borton, A.J., Frederick, J.P., Datto, M.B., Wang, X.F., and Weinstein, R.S. (2001). The loss of Smad3 results in a lower rate of bone formation and osteopenia through dysregulation of osteoblast differentiation and apoptosis. *J Bone Miner Res* 16, 1754-1764.
42. Claro da Silva, T., Hiller, C., Gai, Z., and Kullak-Ublick, G.A. (2016). Vitamin D3 transactivates the zinc and manganese transporter SLC30A10 via the Vitamin D receptor. *J Steroid Biochem Mol Biol* 163, 77-87.
43. O'Seaghdha, C.M., Wu, H., Yang, Q., Kapur, K., Guessous, I., Zuber, A.M., Kottgen, A., Stoudmann, C., Teumer, A., Kutalik, Z., et al. (2013). Meta-analysis of genome-wide association studies identifies six new Loci for serum calcium concentrations. *PLoS Genet* 9, e1003796.
44. Speliotes, E.K., Yerges-Armstrong, L.M., Wu, J., Hernaez, R., Kim, L.J., Palmer, C.D., Gudnason, V., Eiriksdottir, G., Garcia, M.E., Launer, L.J., et al. (2011). Genome-wide association analysis identifies variants associated with nonalcoholic fatty liver disease that have distinct effects on metabolic traits. *PLoS Genet* 7, e1001324.
45. Ousingsawat, J., Wanitchakool, P., Schreiber, R., Wuelling, M., Vortkamp, A., and Kunzelmann, K. (2015). Anoctamin-6 controls bone mineralization by activating the calcium transporter NCX1. *J Biol Chem* 290, 6270-6280.
46. Shy, B.R., Wu, C.I., Khramtsova, G.F., Zhang, J.Y., Olopade, O.I., Goss, K.H., and Merrill, B.J. (2013). Regulation of Tcf7l1 DNA binding and protein stability as principal mechanisms of Wnt/beta-catenin signaling. *Cell Rep* 4, 1-9.
47. Tsutsumi, K., Matsuda, M., Kotani, M., Mizokami, A., Murakami, A., Takahashi, I., Terada, Y., Kanematsu, T., Fukami, K., Takenawa, T., et al. (2011). Involvement of PRIP, phospholipase C-related, but catalytically inactive protein, in bone formation. *J Biol Chem* 286, 31032-31042.
48. Li, Y., and Dudley, A.T. (2009). Noncanonical frizzled signaling regulates cell polarity of growth plate chondrocytes. *Development* 136, 1083-1092.
49. Miclea, R.L., Karperien, M., Bosch, C.A., van der Horst, G., van der Valk, M.A., Kobayashi, T., Kronenberg, H.M., Rawadi, G., Akcakaya, P., Lowik, C.W., et al. (2009). Adenomatous polyposis coli-mediated control of beta-catenin is essential for both chondrogenic and osteogenic differentiation of skeletal precursors. *BMC Dev Biol* 9, 26.
50. Wu, Q.T., Ma, Q.J., He, C.Y., Wang, C.X., Gao, S., Hou, X., and Ma, T.H. (2007). Reduced bone mineral density and bone metabolism in aquaporin-1 knockout mice. *Chem Res Chinese U* 23, 297-299.

51. Zhai, G.J., Teumer, A., Stolk, L., Perry, J.R.B., Vandenput, L., Coviello, A.D., Koster, A., Bell, J.T., Bhasin, S., Eriksson, J., et al. (2011). Eight Common Genetic Variants Associated with Serum DHEAS Levels Suggest a Key Role in Ageing Mechanisms. *Plos Genetics* 7.
52. Moon, S.J., Lim, M.A., Park, J.S., Byun, J.K., Kim, S.M., Park, M.K., Kim, E.K., Moon, Y.M., Min, J.K., Ahn, S.M., et al. (2014). Dual-specificity phosphatase 5 attenuates autoimmune arthritis in mice via reciprocal regulation of the Th17/Treg cell balance and inhibition of osteoclastogenesis. *Arthritis Rheumatol* 66, 3083-3095.
53. Hayer, S., Steiner, G., Gortz, B., Reiter, E., Tohidast-Akrad, M., Amling, M., Hoffmann, O., Redlich, K., Zwerina, J., Skriner, K., et al. (2005). CD44 is a determinant of inflammatory bone loss. *J Exp Med* 201, 903-914.
54. Krishnan, V., Bryant, H.U., and Macdougald, O.A. (2006). Regulation of bone mass by Wnt signaling. *J Clin Invest* 116, 1202-1209.
55. Ryan, Z.C., Craig, T.A., Filoteo, A.G., Westendorf, J.J., Cartwright, E.J., Neyses, L., Strehler, E.E., and Kumar, R. (2015). Deletion of the intestinal plasma membrane calcium pump, isoform 1, Atp2b1, in mice is associated with decreased bone mineral density and impaired responsiveness to 1, 25-dihydroxyvitamin D3. *Biochem Biophys Res Commun* 467, 152-156.
56. Tsukamoto, S., Mizuta, T., Fujimoto, M., Ohte, S., Osawa, K., Miyamoto, A., Yoneyama, K., Murata, E., Machiya, A., Jimi, E., et al. (2014). Smad9 is a new type of transcriptional regulator in bone morphogenetic protein signaling. *Sci Rep* 4, 7596.
57. Miyaura, C., Toda, K., Inada, M., Ohshiba, T., Matsumoto, C., Okada, T., Ito, M., Shizuta, Y., and Ito, A. (2001). Sex- and age-related response to aromatase deficiency in bone. *Biochem Biophys Res Commun* 280, 1062-1068.
58. Kim, K., Kim, J.H., Lee, J., Jin, H.M., Kook, H., Kim, K.K., Lee, S.Y., and Kim, N. (2007). MafB negatively regulates RANKL-mediated osteoclast differentiation. *Blood* 109, 3253-3259.
59. Tamamura, Y., Otani, T., Kanatani, N., Koyama, E., Kitagaki, J., Komori, T., Yamada, Y., Costantini, F., Wakisaka, S., Pacifici, M., et al. (2005). Developmental regulation of Wnt/beta-catenin signals is required for growth plate assembly, cartilage integrity, and endochondral ossification. *J Biol Chem* 280, 19185-19195.
60. Soung do, Y., Kalinowski, J., Baniwal, S.K., Jacome-Galarza, C.E., Frenkel, B., Lorenzo, J., and Drissi, H. (2014). Runx1-mediated regulation of osteoclast differentiation and function. *Mol Endocrinol* 28, 546-553.

61. Li, V., Raouf, A., Kitching, R., and Seth, A. (2004). Ets2 transcription factor inhibits mineralization and affects target gene expression during osteoblast maturation. *In Vivo* 18, 517-524.
62. Tijchon, E., van Ingen Schenau, D., van Opzeeland, F., Tirone, F., Hoogerbrugge, P.M., Van Leeuwen, F.N., and Scheijen, B. (2015). Targeted Deletion of Btg1 and Btg2 Results in Homeotic Transformation of the Axial Skeleton. *PLoS One* 10, e0131481.
63. Koscielny, G., Yaikhom, G., Iyer, V., Meehan, T.F., Morgan, H., Atienza-Herrero, J., Blake, A., Chen, C.K., Easty, R., Di Fenza, A., et al. (2014). The International Mouse Phenotyping Consortium Web Portal, a unified point of access for knockout mice and related phenotyping data. *Nucleic Acids Res* 42, D802-809.
64. Ghebre, M.A., Hart, D.J., Hakim, A.J., Kato, B.S., Thompson, V., Arden, N.K., Spector, T.D., and Zhai, G.J. (2011). Association between DHEAS and Bone Loss in Postmenopausal Women: A 15-Year Longitudinal Population-Based Study. *Calcified Tissue Int* 89, 295-302.
65. Gong, Y., Slee, R.B., Fukai, N., Rawadi, G., Roman-Roman, S., Reginato, A.M., Wang, H., Cundy, T., Glorieux, F.H., Lev, D., et al. (2001). LDL receptor-related protein 5 (LRP5) affects bone accrual and eye development. *Cell* 107, 513-523.
66. Heilmeier, U., Hackl, M., Skalicky, S., Weilner, S., Schroeder, F., Vierlinger, K., Patsch, J.M., Baum, T., Oberbauer, E., Lobach, I., et al. (2016). Serum miRNA Signatures Are Indicative of Skeletal Fractures in Postmenopausal Women With and Without Type 2 Diabetes and Influence Osteogenic and Adipogenic Differentiation of Adipose Tissue-Derived Mesenchymal Stem Cells In Vitro. *J Bone Miner Res* 31, 2173-2192.
67. Sekiya, H., Murakami, T., Saito, A., Hino, S., Tsumagari, K., Ochiai, K., and Imaizumi, K. (2010). Effects of the bisphosphonate risedronate on osteopenia in OASIS-deficient mice. *J Bone Miner Metab* 28, 384-394.
68. Boudin, E., Yorgan, T., Fijalkowski, I., Sonntag, S., Steenackers, E., Hendrickx, G., Peeters, S., Mare, A., Vervae, B., Verhulst, A., et al. (2017). The Lrp4 R1170Q homozygous knock-in mouse recapitulates the bone phenotype of sclerosteosis in humans. *J Bone Miner Res*.
69. Albagha, O.M., Wani, S.E., Visconti, M.R., Alonso, N., Goodman, K., Brandi, M.L., Cundy, T., Chung, P.Y., Dargie, R., Devogelaer, J.P., et al. (2011). Genome-wide association identifies three new susceptibility loci for Paget's disease of bone. *Nat Genet* 43, 685-689.
70. Courant, F., Aksglaede, L., Antignac, J.P., Monteau, F., Sorensen, K., Andersson, A.M., Skakkebaek, N.E., Juul, A., and Bizec, B.L. (2010). Assessment of circulating sex steroid levels in prepubertal

and pubertal boys and girls by a novel ultrasensitive gas chromatography-tandem mass spectrometry method. *J Clin Endocrinol Metab* 95, 82-92.

71. Cao, J., Venton, L., Sakata, T., and Halloran, B.P. (2003). Expression of RANKL and OPG correlates with age-related bone loss in male C57BL/6 mice. *J Bone Miner Res* 18, 270-277.
72. Yang, J., Bakshi, A., Zhu, Z., Hemani, G., Vinkhuyzen, A.A., Lee, S.H., Robinson, M.R., Perry, J.R., Nolte, I.M., van Vliet-Ostaptchouk, J.V., et al. (2015). Genetic variance estimation with imputed variants finds negligible missing heritability for human height and body mass index. *Nat Genet* 47, 1114-1120.

Figure Titles and Legends

Figure 1. Manhattan plot of association statistics ($-\log_{10}(P\text{-values})$) for TB-BMD overall meta-analysis. Each dot represents a SNP and the x-axis indicates its chromosomal position (built 37 NCBI). Red dots represent SNPs at GWS loci that are not within $\pm 500\text{Kb}$ of leading SNPs in previous GWAS with different bone traits. Dashed horizontal red and yellow lines mark the GWS threshold ($P < 5 \times 10^{-8}$) and suggestive threshold ($P < 1 \times 10^{-6}$), respectively. Novel loci in the only-CEU analysis are not shown.

Figure 2. Age dependence of the genetic variant effect in the meta-regression. The panels display leading SNPs from two loci exhibiting significant evidence for age influences. Heterogeneity P-values (P_{het}) are reported for the overall meta-analysis. In the left panels, each circle represents a study subgroup (i.e., study divided in age strata), with the circle size proportional to the inverse variance of the SNP main effect. In the right panels, forest plots display estimates obtained from each age-bin meta-analysis, with the symbol size proportional to the inverse variance of the SNP main effect.

Figure 3. Genetic correlations between TB-BMD and other traits and diseases. Calculation was based on the summary statistics of the only-European meta-analysis ($N=56,284$) and estimated by LD score regression implemented in LDHub. The diagram only show traits whose correlation with TB-BMD was significant ($P < 0.05$).

Figure 4. Depict results for gene-set and cell/tissue enrichment analyses. **Top panel:** 25 Meta gene-sets were defined from similarity clustering of significantly enriched gene sets ($\text{FDR} < 5\%$). Each Meta gene-set was named after one of its member gene sets. The color of the Meta gene-sets represents the P-value of the member set. Interconnection line width represents the Pearson correlation ρ between the gene membership scores for each Meta gene-set ($\rho < 0.3$, no line; $0.3 \leq \rho < 0.5$, narrow width; $0.5 \leq \rho < 0.7$, medium width; $\rho \geq 0.7$, thick width). **Bottom panel:** Bars represent the level of evidence for genes in the associated loci to be expressed in any of the 209 Medical Subject Heading (MeSH) tissue and cell type annotations. Highlighted in orange are these cell/tissue types significantly ($\text{FDR} < 5\%$) enriched for the expression of the genes in the associated loci.

Tables

Table 1. Index SNPs of loci not previously associated with BMD. Variants associated with TB-BMD in the all-ages combined meta-analysis that map outside +/- 500 Kb of known index SNPs of genetic associations with different bone traits. Genomic coordinates are on build 37 of the human genome. Notes refer to annotation based on the closest gene. Associations with Lumbar Spine (LS) and Femoral Neck (FN)-BMD¹⁰. Beta coefficients and allele frequencies (EAF) are reported for the A1 allele

CHR	BP	rsnumber	Locus	A1	A2	EAF	Effect	P	N	annotation	closest gene	Notes	LS-beta	LS-P	FN-beta	FN-P
1	8422676	rs2252865	1p36.23	T	C	0.32	-0.033	4.72E-08	66075	intronic	<i>RERE</i>	Novel biology	-0.019	0.043	-0.025	0.002
1	110475971	rs7548588	1p13.3	T	C	0.61	-0.037	9.29E-09	66240	intergenic	<i>CSF1</i>	Osteoclast differentiation ⁴⁰	-0.030	0.001	-0.022	0.005
1	220038825	rs185048405	1q41	T	C	0.54	0.042	3.07E-09	66540	intronic	<i>SLC30A10</i>	Manganese transport ⁴²	-0.035	0.076	-0.003	0.878
2	27741072	rs780096	2p23.3	C	G	0.44	-0.031	4.58E-08	66578	intronic	<i>GCKR</i>	Calcium regulation ⁴³ , hepatic traits ⁴⁴	-0.014	0.129	-0.017	0.029
2	40630678	rs10490046	2p22.1	A	C	0.76	0.043	1.43E-10	65961	intronic	<i>SLC8A1</i>	Bone mineralization ⁴⁵	0.015	0.162	0.021	0.025
2	68962137	rs10048745	2p13.3	A	G	0.25	-0.039	6.44E-09	66565	5'-UTR	<i>ARHGAP25</i>	Novel biology	-0.050	1.03E-06	-0.036	5.21E-05
2	85484818	rs11904127	2p11.2	A	G	0.55	-0.032	2.65E-08	66561	intronic	<i>TCF7L1</i>	Factors in Wnt signaling ⁴⁶	-0.021	0.023	-0.015	0.054
2	198874006	rs1595824	2q33.1	T	C	0.47	0.034	2.65E-08	60171	intronic	<i>PLCL1</i>	Negative regulation of bone formation ⁴⁷	0.022	0.201	0.052	2.20E-04
2	202799604	rs2350085	2q33.2	T	C	0.87	-0.064	3.80E-14	66412	intergenic	<i>FZD7</i>	Factors in Wnt signaling ⁴⁸	-0.042	0.002	-0.044	1.96E-04
2	234303405	rs838721	2q37.1	A	G	0.44	-0.031	4.48E-09	65516	intronic	<i>DGKD</i>	Calcium regulation ⁴³	-0.016	0.070	-0.014	0.068
5	112221869	rs818427	5q22.2	T	C	0.31	0.034	2.37E-08	66592	intronic	<i>APC</i>	Bone metabolism ⁴⁹	0.004	0.645	0.008	0.327
5	122847622	rs11745493	5q23.2	A	G	0.75	0.044	7.75E-12	66597	promoter	<i>CSNK1G3</i>	Novel Biology	0.010	0.326	0.025	0.005
7	27989403	rs757138	7p15.1	T	G	0.69	-0.035	3.33E-08	66043	intronic	<i>JAZF1</i>	Novel Biology	-0.016	0.126	-0.025	0.004
7	30957702	rs28362721	7p14.3	T	C	0.18	-0.059	6.71E-14	66274	intronic	<i>AQP1</i>	Bone metabolism ⁵⁰	-0.037	0.002	-0.049	1.39E-06
7	50901491	rs1548607	7p12.1	A	G	0.69	0.036	4.18E-08	66564	intergenic	<i>GRB10</i>	Novel biology	0.034	5.59E-04	0.005	0.517
7	99130834	rs34670419	7q22.1	T	G	0.04	-0.088	1.09E-08	66336	3'-UTR	<i>ZKSCAN5</i>	DHEAS and aging mechanisms ⁵¹	-0.127	9.28E-08	-0.080	8.19E-05
10	112245400	rs73349318	10q25.2	A	T	0.87	-0.047	2.68E-08	66341	intronic	<i>DUSP5</i>	Osteoclast differentiation ⁵²	-0.042	0.001	-0.051	8.76E-06
10	124015986	rs10788264	10q26.13	A	G	0.48	-0.034	2.61E-09	66565	intergenic	<i>TACC2</i>	Novel Biology	-0.030	9.64E-04	-0.029	1.29E-04
11	242859	rs55781332	11p15.5	A	G	0.78	-0.055	8.07E-16	66198	intronic	<i>PSMD13</i>	Novel Biology	-0.046	1.76E-05	-0.026	0.005
11	35083633	rs2553773	11p13	C	G	0.41	-0.037	1.49E-10	66619	intergenic	<i>CD44</i>	Osteoclast activity ⁵³	-0.015	0.101	-0.015	0.054
11	35981346	rs113964474*	11p.13*	A	G	0.03	0.485	1.41E-08	6748	intronic	<i>LDLRAD3</i>	Novel Biology
11	69299537	rs4980659	11q13.3	C	G	0.52	0.033	1.16E-08	66537	intergenic	<i>CCND1</i>	Target of Wnt signalling ⁵⁴	0.039	1.58E-05	0.023	0.003
11	121913230	rs725670	11q24.1	A	G	0.38	-0.032	3.61E-08	66565	intergenic	<i>BLID</i>	Novel Biology	-0.020	0.028	-0.011	0.172
12	90334829	rs10777212	12q21.33	T	G	0.35	0.045	5.05E-14	66619	intergenic	<i>ATP2B1</i>	Calcium absorption ⁵⁵	0.028	0.003	0.021	0.010
12	116555786	rs73200209**	12q24.21	A	T	0.80	0.045	2.51E-08	51240	intronic	<i>MED13L</i>	Novel biology	0.030	0.167	0.036	0.044
13	37487021	rs556429	13q13.3	A	C	0.23	0.039	1.46E-08	66504	intronic	<i>SMAD9</i>	Osteoblast differentiation ⁵⁶	0.023	0.027	0.013	0.135
15	38340874	rs12442242	15q14	A	G	0.85	-0.051	4.94E-10	66403	intergenic	<i>TMC05A</i>	Novel Biology	-0.046	3.03E-04	-0.047	2.26E-05
15	51537806	rs2414098	15q21.2	T	C	0.39	-0.033	1.99E-08	66562	intronic	<i>CYP19A1</i>	Estrogen byosynthesis ⁵⁷	-0.034	0.007	-0.038	0.001
15	67420680	rs1545161	15q22.33	A	G	0.56	0.041	1.06E-12	66004	intronic	<i>SMAD3</i>	Osteoblast differentiation ⁴¹	0.034	1.27E-04	0.035	5.78E-06
17	17804725	rs8070128	17p11.2	T	C	0.58	-0.039	1.98E-11	66625	intronic	<i>TOM1L2</i>	Novel biology	-0.033	4.80E-04	-0.015	0.052
17	63771079	rs9972944	17q24.1	A	G	0.41	0.036	6.87E-10	66595	intronic	<i>CEP112</i>	Novel Biology	0.028	0.003	0.004	0.576
19	31654615	rs6510186***	19q12	T	C	0.26	0.068	3.11E-08	18782	intergenic	<i>TSHZ3</i>	Novel Biology	0.004	0.713	0.006	0.492

20	39103882	rs6029130	20q12	T	C	0.30	0.035	3.50E-08	66497	intergenic	<i>MAFB</i>	Osteoclast differentiation ⁵⁸	0.027	0.007	0.015	0.083
21	28773868	rs1452102	21q21.3	T	G	0.59	-0.035	1.74E-09	66489	intergenic	<i>ADAMTS5</i>	Endochondral Ossification ⁵⁹	-0.029	0.001	-0.015	0.056
21	36970350	rs9976876	21q22.12	T	G	0.45	-0.038	8.01E-11	66514	intronic	<i>RUNX1</i>	Osteoclast differentiation ⁶⁰	-0.019	0.031	-0.016	0.041
21	40350744	rs11910328	21q22.2	A	G	0.84	-0.043	2.99E-08	66298	intergenic	<i>ETS2</i>	Osteoblast maturation ⁶¹	-0.028	0.020	-0.028	0.007

* Monomorphic in European cohorts. ** Reported statistics from the in the meta-analysis of European populations. *** Reported statistics from the meta-analysis in the 30-45 age-strata.